

Hierarchical renormalization group fixed points

A. Pordt*

Institut für Theoretische Physik I, Universität Münster,
Wilhelm-Klemm-Str. 9, D-48149 Münster, Germany

February 1, 2008

Abstract

Hierarchical renormalization group transformations are related to non-associative algebras. Non-trivial infrared fixed points are shown to be solutions of polynomial equations. At the example of a scalar model in $d(\geq 2)$ dimensions some methods for the solution of these algebraic equations are presented.

1 Introduction

Wilson's renormalization group (RG) approach [1] is an important tool for the investigation of quantum field theories. The study of the ultraviolet (UV) and infrared (IR) limits correspond to the study of field theories near fixed points of a renormalization group transformation. A trivial fixed point (UV fixed point) corresponds to asymptotically free theories. For these models standard perturbation methods are sufficient. There are other non-trivial RG fixed points (IR fixed points) where a small parameter is lacking. The goal of this paper is to present non-perturbative methods for analyzing models at IR fixed points.

Investigation of quantum field theories at infrared fixed points requires non-perturbative methods. In this paper we want to consider hierarchical RG fixed points.

The hierarchical RG transformation (HRGT) for a scalar field theory is defined in the following. Let $Z : \mathbf{R} \rightarrow \mathbf{R}$ be a (Gaussian-measurable) function (which corresponds to the generating functional of free propagator-amputated Greens functions). Z is also called *partition function*. Application of the HRGT $\mathcal{RG}$ to the partition function Z yields the *effective partition function* $\mathcal{RG}(Z) : \mathbf{R} \rightarrow \mathbf{R}$ defined by

$$\mathcal{RG}(Z)(\Psi) := \int d\mu_\gamma(\Phi) \left[Z(\Phi + L^{1-\frac{d}{2}}\Psi) \right]^{L^d}, \quad (1)$$

*Work supported by the Deutsche Forschungsgemeinschaft under Grant Wi 1280/2-1

where $\Psi \in \mathbf{R}$, $L \in \{2, 3, \dots\}$, $\gamma > 0$ and the Gaussian measure is defined by

$$\int d\mu_\gamma(\Phi) := \int_{-\infty}^{\infty} d\Phi \exp\left\{-\frac{\Phi^2}{2\gamma}\right\}. \quad (2)$$

The factor L corresponds to the block size of a block spin RG transformation and γ corresponds to the fluctuation propagator. The generalization to N -component models is straightforward.

We want to find even fixed points Z^* of the HRGT $Z^* = \mathcal{RG}(Z^*)$. Excluding $Z^* = 0$ or ∞ , we immediately find the following trivial fixed points. There are the UV fixed point $Z_{UV}(\Psi) = 1$ and the high-temperature fixed point $Z_{HT}(\Psi) = L^{\frac{1}{L^d-1}} \exp\left\{-\frac{L^2-1}{2\gamma L^d} \Psi^2\right\}$. For $d \geq 4$ there are only these trivial fixed points. In $2 \leq d < 4$ dimensions it turns out that there exists also other non-trivial fixed points which are called infrared fixed points. For a further investigation one introduces the hierarchical RG algebra [3].

2 Hierarchical renormalization group algebras

Let $P(\Phi)$ be a polynomial in Φ . Define normal-ordering (Wick-ordering) of $P(\Phi)$ by

$$:P(\Phi):_\gamma := \exp\left\{-\frac{\gamma}{2} \frac{\partial^2}{\partial \Phi^2}\right\} P(\Phi). \quad (3)$$

Introduce structure coefficients $\mathcal{C}_l^{mn}$ by the fusion relations

$$:\Phi^{2m}:_\gamma \cdot : \Phi^{2n}:_\gamma = \quad (4)$$

$$\sum_{l: |m-n| \leq l \leq m+n} \mathcal{C}_l^{mn} : \Phi^{2l}:_\gamma. \quad (5)$$

Let $\{e_m, m = 0, 1, 2, \dots\}$ be the canonical basis of the vector space $\mathbf{R}^\infty$. Define a $\times$ -product on $\mathbf{R}^\infty$ by

$$e_m \times e_n := \sum_{l: |m-n| \leq l \leq m+n} \mathcal{C}_l^{mn} e_l. \quad (6)$$

Then $(\mathbf{R}^\infty, \times)$ is a commutative and associative algebra with unit element e_0 .

We will identify partition functions $Z : \mathbf{R} \rightarrow \mathbf{R}$ with elements $z = (z_0, z_1, z_2, \dots) = \sum_{n=0}^{\infty} z_n e_n$ of the algebra $\mathbf{R}^\infty$ by

$$Z(\Phi) = \sum_{n=0}^{\infty} \frac{z_n}{\gamma^n} : (\Phi^2)^n :_\gamma. \quad (7)$$

For $z \in \mathbf{R}^\infty$ and $\beta \in \mathbf{R}$ define the scaling relation $\mathcal{S}_\beta : \mathbf{R}^\infty \rightarrow \mathbf{R}^\infty$ by $\mathcal{S}_\beta(z)_l := \beta^{2l} z_l$. Then one can show that the HRGT eq. (1) – where γ is replaced by $(1 - \beta^2)\gamma$ – is equivalent to

$$\mathcal{RG}(z) = \mathcal{S}_\beta(\underbrace{z \times \dots \times z}_{L^d \text{ factors}}), \quad (8)$$

where $\beta := L^{1-\frac{d}{2}}$. Let us remark that this relation is only helpful if $|\beta| < 1$. This seems to exclude the case $d = 2$. We may avoid this problem by a transformation of the HRGT eq. (1). Define a new partition function F by $F := Z/Z_{HT}$. Then the HRGT of F has the same form as the HRGT eq. (1) of Z . Only the coefficient γ has to be replaced by $L^{-2}\gamma$ and the scaling parameter β by $L^{-2}\beta = L^{-1-\frac{d}{2}} < 1$. This method is called “extraction of the high-temperature fixed point”.

For the simplest choice $L^d = 2$ the fixed points z^* are solutions of the quadratic equation $z = z \times_\beta z := \mathcal{S}_\beta(z \times z)$. The product $\times_\beta$, for $\beta \neq 1$, defines a commutative and non-associative algebra $(\mathbf{R}^\infty, \times_\beta)$ containing no unit element. The algebra $(\mathbf{R}^\infty, \times_\beta)$ is called a *hierarchical renormalization group algebra* (HRGA). By change of coordinates there are also other HRGA’s possible. For example, identify the partition functions $Z : \mathbf{R} \rightarrow \mathbf{R}$ with elements $\tilde{z} = (\tilde{z}_0, \tilde{z}_1, \tilde{z}_2, \dots) = \sum_{n=0}^\infty \tilde{z}_n e_n$ of the algebra $\mathbf{R}^\infty$ by series expansion

$$Z(\Phi) = \sum_{n=0}^\infty \frac{\tilde{z}_n}{\gamma^n} (\Phi^2)^n. \quad (9)$$

Then one can show that the HRGT eq. (1) is equivalent to

$$\mathcal{RG}(\tilde{z}) = \mathcal{S}_\beta(\tilde{z} \star \tilde{z}) = \tilde{z} \star_\beta \tilde{z}, \quad (10)$$

where the $\star_\beta$ -product defines the non-associative algebra $(\mathbf{R}^\infty, \star_\beta)$ and the corresponding structure coefficients $\mathcal{S}_l^{mn}$ can be easily computed [2, 3]. Let us remark that this product cannot be used for cases $L^d \neq 2$, since $(\mathbf{R}^\infty, \star_\beta)$ is also non-associative for $\beta = 1$.

For periodic fixed points in $d = 2$ dimensions one represents the partition function Z with period $T = \frac{\sqrt{\gamma}}{q}$ by

$$Z(\Phi) = \sum_{m0=-\infty}^\infty z_m \exp\{2\pi i m \frac{q}{\sqrt{\gamma}} \Phi\} \quad (11)$$

Define a $\circ_q$ -product by

$$e_m \circ_q e_n := \exp\{-2\pi^2 q^2 (m+n)^2\} e_{m+n}. \quad (12)$$

Then the HRGT eq. (1), for $d = 2$ and $L^2 = 2$, is equivalent to

$$\mathcal{RG}(z) = z \circ_q z \quad (13)$$

The HRGA $(\mathbf{R}^\infty, \circ_q)$ is non-associative for $q \neq 0$.

2.1 Norms

For estimations and convergence proofs suitable definitions of norms for HRGA’s are required. Equipped with these norms the algebras $\mathbf{R}^\infty$ become Banach-algebras.

The norms $\|\cdot\|_\rho$, defined below, depend on a parameter ρ . If the parameter ρ is large enough, $\rho \geq \rho^*$, the norms become algebra-norms, i. e. they obey

$$\|a \bullet b\|_\rho \leq \|a\|_\rho \cdot \|b\|_\rho, \quad (14)$$

where the $\bullet$ -product defines a HRGA. For example, Koch and Wittwer [4] used the following norm for their proof of a non-trivial fixed point in $d = 3$ dimensions for the case $L^d = 2$ and extracting the high-temperature fixed point ($2\beta^2 < 1$)

$$\|z\|_\rho^{(1)} := \sum_{n=0}^{\infty} \sqrt{(2n)!} |z_n| \rho^n. \quad (15)$$

For the use of estimating coefficients in the ϵ -expansion the following sup-norm is useful [3] ($2\beta^2 > 1$)

$$\|z\|_\rho^{(\infty)} := \sup_n (n! |z_n| \rho^n). \quad (16)$$

In constructive field theory norms are typically defined by a large field regulator $e^{\rho\Phi^2}$

$$\|z\|_\rho^{(c)} := \sup_{\Phi \in \mathbf{R}} |e^{\rho\Phi^2} Z(\Phi)|. \quad (17)$$

3 Methods and Results

There are several methods to solve the fixed point equation. Beside the method of ϵ -expansion all methods presented here deliver convergent expansions for the fixed points.

The first and most standard one is to iterate the RGT. Define a sequence of partition functions by $z^{(n+1)} := \mathcal{RG}(z^{(n)})$. One has to choose the start of the RGT $z^{(0)}$ such that $\lim_{n \rightarrow \infty} z^{(n)} = z^*$ exists. Then z^* is a fixed point.

A second standard method is the ϵ -expansion [5, 2]. Take the trivial UV fixed point $z_{UV} = e_0$ in d_* dimensions. Then one considers the fixed point equation in $d = d_* - \epsilon$ dimensions. The fixed point z^* and the scaling factor β^{2l} is written as a power series in ϵ . Insertion of these series in the fixed point equation $z = z \times_\beta z$ and comparison of the coefficients yields recursive equations for the coefficients of the fixed point in the case of critical dimensions $d_* = d_l := \frac{2l}{l-1}$, $l \in \{2, 3, \dots\}$. $\ln Z^*(\Phi)$ has the form of an l -well.

That d_l are the critical dimensions can be seen in the following way. We suppose that for small ϵ the fixed point $z = e_0 + r$ is near by the UV fixed point e_0 , i. e. r is small. The fixed point equation in terms of r is $r = 2e_0 \times_\beta r + r \times_\beta r$. In lowest order r should obey $r \in \ker(id - 2e_0 \times_\beta)$. The linear operator $(id - 2e_0 \times_\beta) = \text{diag}(1 - 2\beta^{2l}, l = 0, 1, \dots)$ has only a non-trivial kernel if there exists l such that $1 = 2\beta^{2l}$. This is the case for the critical dimensions $d = d_l$.

A third method used by Koch and Wittwer [4] is the method of exact beta-functions. For that, one defines a projection operator P on $\mathbf{R}^\infty$ such that

$$z^{rel} = z^{rel}(\gamma_0, \dots, \gamma_N) = P(z) \quad (18)$$

depends only on a finite number of parameters $\gamma_0, \dots, \gamma_N$ and such that the irrelevant part of z defined by $r := (1 - P)(z)$ can be controlled by standard fixed point theorems. The irrelevant fixed point $r^* = r^*(\gamma)$ is defined by

$$r^* = H_\gamma(r^*) := (1 - P)\mathcal{RG}(z^{rel}(\gamma) + r^*). \quad (19)$$

The exact beta-function $B : \mathbf{R}^{N+1} \rightarrow \mathbf{R}^{N+1}$ is defined by

$$z^{rel}(B(\gamma_0, \dots, \gamma_N)) = \quad (20)$$

$$P\mathcal{RG}(z^{rel}(\gamma_0, \dots, \gamma_N) + r^*(\gamma_0, \dots, \gamma_N)). \quad (21)$$

The fixed points $z^* = \mathcal{RG}(z^*)$ are given by $z^* = z^{rel}(\gamma^*) + r^*(\gamma^*)$, where γ^* is the fixed point of the exact beta-function B , $B(\gamma^*) = \gamma^*$. For the HRGT one can use the projection operator

$$P(z) := (z_0, \dots, z_N, 0, 0, \dots) \quad (22)$$

and the beta-function technique works for $N \geq N_0 = O(1)(= 7)$.

A fourth method is to replace the fixed point equation by a minimization-problem. Define an energy-function $E(z)$ for $z \in \mathbf{R}^\infty$ by

$$E(z) := \sum_{l=0}^{\infty} (z_l - (z \times_\beta z)_l)^2 \geq 0. \quad (23)$$

Then $z = z \times_\beta z$ is equivalent to $E(z) = 0$.

A fifth method is to solve the equation $z \times_\beta z - z = 0$ directly by Newton's method [2].

A sixth method is to truncate the infinite system of quadratic equations by setting $z_l = 0$ if $l > N$ and solve the finite system directly by means of Groebner bases.

For convergence proofs and numerical calculations one has to truncate the infinite system. One can show that there exists a solution of the truncated fixed point equation $z^{(N)}$ such that the limit $\lim_{N \rightarrow \infty} z^{(N)} = z^*$ exists.

References

- [1] K. G. Wilson, Physica 73 (1974) 119; K. G. Wilson, J. Kogut, Phys. Rep. C12 (1974) 12.
- [2] K. Pinn, A. Pordt, C. Wiecekowsky, Journ. Stat. Phys. 77 (1994) 977.
- [3] A. Pordt, C. Wiecekowsky, MS-TPI-94-04.
- [4] H. Koch, P. Wittwer, Commun. Math. Phys. 106 (1986) 495; Commun. Math. Phys. 138 (1991) 537.
- [5] P. Collet, J.-P. Eckmann, Commun. Math. Phys. 55 (1977) 67.